

Distance Between Point and Line, Ray, or Line Segment

David Eberly
Geometric Tools, LLC
<http://www.geometrictools.com/>
Copyright © 1998-2016. All Rights Reserved.

Created: March 2, 1999
Last Modified: March 1, 2008

Contents

1 Discussion

2

1 Discussion

The following construction applies in any dimension, not just in 3D. Let the test point be \mathbf{P} . A line is parameterized as $\mathbf{L}(t) = \mathbf{B} + t\mathbf{M}$ where \mathbf{B} is a point on the line, \mathbf{M} is the line direction, and $t \in \mathbb{R}$. A *ray* is of the same form but with restriction $t \geq 0$. A *line segment* is restricted even further with $t \in [0, 1]$. The end points of the line segment are \mathbf{B} and $\mathbf{B} + \mathbf{M}$.

The closest point on the line to \mathbf{P} is the projection of \mathbf{P} onto the line, $\mathbf{Q} = \mathbf{B} + t_0\mathbf{M}$, where

$$t_0 = \frac{\mathbf{M} \cdot (\mathbf{P} - \mathbf{B})}{\mathbf{M} \cdot \mathbf{M}}.$$

The distance from \mathbf{P} to the line is

$$D = |\mathbf{P} - (\mathbf{B} + t_0\mathbf{M})|.$$

If $t_0 \leq 0$, then the closest point on the ray to \mathbf{P} is \mathbf{B} . For $t_0 > 0$, the projection $\mathbf{B} + t_0\mathbf{M}$ is the closest point. The distance from \mathbf{P} to the ray is

$$D = \left\{ \begin{array}{ll} |\mathbf{P} - \mathbf{B}|, & t_0 \leq 0 \\ |\mathbf{P} - (\mathbf{B} + t_0\mathbf{M})|, & t_0 > 0 \end{array} \right\}.$$

Finally, if $t_0 > 1$, then the closest point on the line segment to \mathbf{P} is $\mathbf{B} + \mathbf{M}$. The distance from \mathbf{P} to the line segment is

$$D = \left\{ \begin{array}{ll} |\mathbf{P} - \mathbf{B}|, & t_0 \leq 0 \\ |\mathbf{P} - (\mathbf{B} + t_0\mathbf{M})|, & 0 < t_0 < 1 \\ |\mathbf{P} - (\mathbf{B} + \mathbf{M})|, & t_0 \geq 1 \end{array} \right\}.$$

The division by $\mathbf{M} \cdot \mathbf{M}$ is the most expensive algebraic operation. The implementation should defer the division as late as possible.